



CN-China: Revised runoff curve number by using rainfall-runoff events data in China

Huishu Lian^{a, b}, Haw Yen^c, Jr-Chuan Huang^d, Qingyu Feng^e, Lihuan Qin^a, Muhammad Amjad Bashir^a, Shuxia Wu^a, A-Xing Zhu^{f, g}, Jiafa Luo^h, Hongjie Diⁱ, Qiuliang Lei^{a, *}, Hongbin Liu^{a, **}

^a Key Laboratory of Nonpoint Source Pollution Control, Ministry of Agriculture, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing, 10081, China

^b School of Environmental Science and Engineering, Southern University of Science and Technology, Shenzhen, Guangdong, 518055, China

^c Blackland Research and Extension Center, Texas A&M Agrilife Research, Texas A&M University, Temple, TX, 76502, USA

^d Department of Geography, National Taiwan University, Taipei, 10617, Taiwan

^e Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China

^f Key Laboratory of Virtual Geographic Environment, Nanjing Normal University, Ministry of Education, Nanjing, 210023, China

^g Department of Geography, University of Wisconsin-Madison, Madison, WI, 53706, USA

^h AgResearch Limited, Ruakura Research Centre, Hamilton, 3240, New Zealand

ⁱ Centre for Soil and Environmental Research, Lincoln University, Lincoln, Christchurch, 7647, New Zealand

ARTICLE INFO

Article history:

Received 22 September 2019

Received in revised form

2 March 2020

Accepted 28 March 2020

Available online 2 April 2020

Keywords:

Hydrology

Curve number

Surface runoff estimation

Modeling

ABSTRACT

The curve number (CN) method developed by the United States Department of Agriculture (USDA) in 1954 is the most common adopted method to estimate surface runoff. For years, applicability of the CN method is a conundrum when implementing to other countries. Specifically, countries with more complex natural environment may require more dedicated adjustments. Therefore, the current CN look-up table provided by USDA might not be appropriate and could be questionable to be applied directly to regions elsewhere. Some studies have been conducted to modify CN values according to specified natural characteristics in scattered regions of mainland China. However, an integral and representative work is still not available to address potential concerns in general matters. In this study, a large set of rainfall-runoff monitoring data were collected to adjust CN values in 55 study sites across China. The results showed that the revised CN values are largely different from CN look-up table provided by USDA, which would lead to huge errors in runoff estimation. In this study, the revised CN (dubbed CN-China) provides better reference guidelines that are suitable for most natural conditions in China. In addition, scientists and engineers from other parts of the world can take advantage of the proposed work to enhance the quality of future programs related to surface runoff estimation.

© 2020 Elsevier Ltd. All rights reserved.

1. Introduction

The high-density construction of meteorological stations around the world makes it easy to acquire historical and real-time rainfall data, but there are few stations for runoff monitoring. Runoff monitoring is often carried out for short-term and limited area in order to accomplish specific research projects (Fauvel et al., 2016; Gwynne and Glover, 1961; Hvitvedjacobsen and Yousef, 1988; Kim and Sansalone,

2008). Surface runoff estimation serves as the most important role in hydrology related research (Hawkins, 1993; Jiang et al., 2012; Kim and Sansalone, 2008; Kirchner et al., 2000; Muche et al., 2019; Steenhuis et al., 1995; Tyagi et al., 2008; Wang and Wang, 2018). Fortunately, the relationship between runoff and rainfall allows us to use mathematical methods to estimate runoff based on readily available rainfall data (Clyde and Work, 1943; Guo et al., 2017). The Green-Ampt infiltration curve (Freyberg et al., 1980; Li et al., 2015; Stewart, 2018), the Philip infiltration curve (Triadis and Broadbridge, 2012), and the Horton infiltration curve methods are all for calculating runoff (Esen, 1987; Grimaldi et al., 2013). Applications of these methods are very limited because they require many parameters and detailed soil

* Corresponding author.

** Corresponding author.

E-mail addresses: leiqliang@caas.cn (Q. Lei), liuhongbin@caas.cn (H. Liu).